

Toda Lattice and Tomimatsu-Sato Solutions

TAKESHI FUKUYAMA[†], KIYOSHI KAMIMURA[‡] AND SONGJU YU[†]

[†]*Department of Physics, Ritsumeikan University*

Kusatsu, Shiga 525 JAPAN

[‡]*Department of Physics, Toho University*

Funabashi, 274 Japan

e-mails: FUKUYAMA @ JPNYITP, KAMIMURA @ JPNYITP

Abstract

We discuss an analytic proof of a conjecture (Nakamura) that solutions of Toda molecule equation give those of Ernst equation giving Tomimatsu-Sato solutions of Einstein equation. Using Pfaffian identities it is shown for Weyl solutions completely and for generic cases partially.

April 1995
Rits-TH-9511
TOHO-FP-9551

Many of integrable non linear models are obtained from (anti) self-dual Yang-Mills equations by reduction [1](Ward's conjecture). Toda Lattice and Ernst equations are such examples of integrable systems. However the relationship between these integrable systems is not so simple. It is not affirmative that the system of Ernst equation belongs to the conventional category of integrable systems. For instance, Lax pair for the Ernst equation is not exactly the same type as those in typical soliton theories[2].

On the other hand, "*direct method*" developed by Hirota[3] has shown to be a very powerful tool in soliton theories. One of the advantages of the direct method amongst others is that the proofs of soliton solutions are reduced to Pfaffian identities. The mathematical implication of the direct method was revealed by Sato[4]. Both the methods by Hirota and Sato seem to be quite useful in soliton theories. However both theories have been seldom applied to the system of Ernst equation except in very few articles. In applying them to the system of Ernst equation we expect to find the universal characters of soliton theories also in the Ernst equation. This is one of motivations of this article.

Recently Nakamura found an important relation between Tomimatsu-Sato solutions (hereafter we refer it to TS) and Toda lattice solutions [5]. He has shown that a series of TS solutions with deformation parameters n are obtained as n -th lattice site amplitudes in a special case of general solutions of Toda molecule equation (Nakamura's conjecture). He has shown this proposition for small n cases explicitly. The aim of this article is to assure the conjecture analytically and to take any step towards the scheme mentioned above.

We begin with discussing the Toda molecule equation describing a semi-infinite lattice in two dimensions. It is expressed in terms of the Hirota's bilinear forms [3] as

$$(D_X^2 - D_Y^2)\tau_n \cdot \tau_n - 2\tau_{n+1}\tau_{n-1} = 0, \quad (1)$$

where the Hirota derivative is defined by $D(f \cdot g) = (\partial f)g - f(\partial g)$. n 's are positive integers and the boundary condition is $\tau_0 = 1$ corresponding to the semi-infinite lattice. The general solution of Eq.(1) is expressed in a form of two-directional Wronskian [6]

$$\tau_n = \det \begin{pmatrix} \psi & L_- \psi & \dots & L_-^{n-1} \psi \\ L_+ \psi & L_+ L_- \psi & \dots & L_+ L_-^{n-1} \psi \\ \vdots & \vdots & \vdots & \vdots \\ L_+^{n-1} \psi & L_+^{n-1} L_- \psi & \dots & L_+^{n-1} L_-^{n-1} \psi \end{pmatrix}. \quad (2)$$

Here ψ is an arbitrary function and $L_{\pm} \equiv \frac{\partial}{\partial X} \pm \frac{\partial}{\partial Y}$. We first show how it appears as a result of a Pfaffian identity for the help of later discussions, though it is a known fact.

We introduce $\mathcal{D}$ as a determinant of $(n+1) \times (n+1)$ matrix τ_{n+1} ;

$$\mathcal{D} \equiv \tau_{n+1}. \quad (3)$$

The minor $\mathcal{D}\begin{bmatrix} i \\ j \end{bmatrix}$ is defined by deleting the i -th row and the j -th column from $\mathcal{D}$.

Similarly $\mathcal{D}\begin{bmatrix} i,k \\ j,l \end{bmatrix}$ is defined by deleting the i and k -th rows and the j and l -th columns from $\mathcal{D}$ and so on. The Toda molecule equation (1) is now expressed as

$$\mathcal{D}\begin{bmatrix} n \\ n \end{bmatrix} \mathcal{D}\begin{bmatrix} n+1 \\ n+1 \end{bmatrix} - \mathcal{D}\begin{bmatrix} n+1 \\ n \end{bmatrix} \mathcal{D}\begin{bmatrix} n \\ n+1 \end{bmatrix} - \mathcal{D}\mathcal{D}\begin{bmatrix} n, n+1 \\ n, n+1 \end{bmatrix} = 0. \quad (4)$$

It holds since it is nothing but the Jacobi's (Sylvester's) formula for matrix minors, which is one of Pfaffian identities. Thus Toda lattice equation has been reduced to Pfaffian identity in direct method. It is also the case in the Einstein equation as will be shown in the following.

The Nakamura's conjecture on the Tomimatsu-Sato solutions consists of two ingredients [5];

(i) Ernst equation for axially symmetric metric of Einstein equation is

$$(\xi\xi^* - 1)\nabla^2\xi - 2\xi^*\nabla\xi \cdot \nabla\xi = 0. \quad (5)$$

Setting $\xi_n = \frac{g_n}{f_n}$, Eq.(5) has a decomposition into two sets [7]

$$D_x(g_n \cdot f_n - g_n^* \cdot f_n^*) = 0, \quad (6)$$

$$D_y(g_n \cdot f_n + g_n^* \cdot f_n^*) = 0, \quad (7)$$

and

$$F(g_n^* \cdot f_n) = 0, \quad (8)$$

$$F(g_n^* \cdot g_n + f_n^* \cdot f_n) = 0. \quad (9)$$

Here the bi-linear operator F is

$$F = (x^2 - 1)D_x^2 + 2x\partial_x + (y^2 - 1)D_y^2 + 2y\partial_y + c_n \quad (10)$$

and x and y are usual prolated spheroidal coordinates.

(ii) From the solutions of the Toda molecule equation τ_n a set of TS solutions are obtained as

$$g_n = \tau_n = \mathcal{D}\begin{bmatrix} n+1 \\ n+1 \end{bmatrix}, \quad f_n = \tau_{n-1} \big|_{\psi \rightarrow L_+ L_- \psi} = \mathcal{D}\begin{bmatrix} 1, n+1 \\ 1, n+1 \end{bmatrix} \quad (11)$$

by following choices of the arbitrary function ψ and the constant c_n

$$\psi = px - iqy, \quad p^2 + q^2 = 1, \quad c_n = -2n^2. \quad (12)$$

x and X are related by $\partial_X = (x^2 - 1)\partial_x$, and y and Y are related by $\partial_Y = (y^2 - 1)\partial_y$.

It is very suggestive that $\xi_n = \frac{q_n}{f_n}$ in Eq.(11) resembles with the solution W_n of Sato equation[4]. Our plan is to show the second statement (ii) analytically making use of Pfaffian identities. The first set Eqs.(6) and (7) are proved in the generic case while the second set Eqs.(8) and (9) are shown for a restricted case of $q = 0$.

In order to prove the first set of equations Eqs.(6) and (7) we start to redefine $\mathcal{D}$ introduced in Eq.(3). It is fit for treating f_n , g_n and their complex conjugate f_n^* , g_n^* on equal footing. We first introduce complex conjugation operator C ,

$$C\psi = \psi^* = px + ixy, \quad C^2 = \mathbf{1}. \quad (13)$$

We define a new determinant $\mathcal{D}$ of $(n+2) \times (n+2)$ matrix as follows,

$$\mathcal{D} = \det \begin{pmatrix} 0 & C\psi & L_+ C^2 \psi & \dots & L_+^n C^{n+1} \psi \\ \psi & L_+ C \psi & L_+^2 C^2 \psi & \dots & L_+^{n+1} C^{n+1} \psi \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ L_+^n \psi & L_+^{n+1} C \psi & L_+^{n+2} C^2 \psi & \dots & L_+^{2n+1} C^{n+1} \psi \end{pmatrix}. \quad (14)$$

g_n and f_n in Eq.(11) and their conjugate are expressed as minors of $\mathcal{D}$;

$$g_n = \mathcal{D} \begin{bmatrix} 1, n+2 \\ n+1, n+2 \end{bmatrix}, \quad f_n = \mathcal{D} \begin{bmatrix} 1, 2, n+2 \\ 1, n+1, n+2 \end{bmatrix} \quad (15)$$

and

$$g_n^* = \mathcal{D} \begin{bmatrix} n+1, n+2 \\ 1, n+2 \end{bmatrix}, \quad f_n^* = \mathcal{D} \begin{bmatrix} 1, n+1, n+2 \\ 1, 2, n+2 \end{bmatrix}. \quad (16)$$

From the Wronskian structure of $\mathcal{D}$ in Eq.(14), the derivatives of f_n and g_n 's are also expressed as minors

$$\begin{aligned} L_+ g_n &= \mathcal{D} \begin{bmatrix} 1, n+1 \\ n+1, n+2 \end{bmatrix}, & L_+ f_n &= \mathcal{D} \begin{bmatrix} 1, 2, n+1 \\ 1, n+1, n+2 \end{bmatrix}, \\ L_+ g_n^* &= \mathcal{D} \begin{bmatrix} n, n+2 \\ 1, n+2 \end{bmatrix}, & L_+ f_n^* &= \mathcal{D} \begin{bmatrix} 1, n, n+2 \\ 1, 2, n+2 \end{bmatrix} \end{aligned} \quad (17)$$

and

$$\begin{aligned} L_- g_n &= \mathcal{D} \begin{bmatrix} 1, n+2 \\ n, n+2 \end{bmatrix}, & L_- f_n &= \mathcal{D} \begin{bmatrix} 1, 2, n+2 \\ 1, n, n+2 \end{bmatrix}, \\ L_- g_n^* &= \mathcal{D} \begin{bmatrix} n+1, n+2 \\ 1, n+1 \end{bmatrix}, & L_- f_n^* &= \mathcal{D} \begin{bmatrix} 1, n+1, n+2 \\ 1, 2, n+1 \end{bmatrix}. \end{aligned} \quad (18)$$

We will prove that f_n and g_n given in Eq.(11) satisfy Eqs.(6) and (7). In showing it the explicit form of ψ in Eq.(12) is not required. In terms of $L_\pm$, Eq.(6) $\pm$ Eq.(7) are given as

$$g_n L_+ f_n - (L_+ g_n) f_n - g_n^* L_- f_n^* + (L_- g_n^*) f_n^* = 0, \quad (19)$$

$$g_n L_- f_n - (L_- g_n) f_n - g_n^* L_+ f_n^* + (L_+ g_n^*) f_n^* = 0. \quad (20)$$

Using Eqs.(17) and (18), Eq.(19) becomes

$$\begin{aligned}
& \mathcal{D} \begin{bmatrix} 1, n+2 \\ n+1, n+2 \end{bmatrix} \mathcal{D} \begin{bmatrix} 1, 2, n+1 \\ 1, n+1, n+2 \end{bmatrix} \\
& - \mathcal{D} \begin{bmatrix} 1, n+1 \\ n+1, n+2 \end{bmatrix} \mathcal{D} \begin{bmatrix} 1, 2, n+2 \\ 1, n+1, n+2 \end{bmatrix} \\
& - \mathcal{D} \begin{bmatrix} n+1, n+2 \\ 1, n+2 \end{bmatrix} \mathcal{D} \begin{bmatrix} 1, n+1, n+2 \\ 1, 2, n+1 \end{bmatrix} \\
& + \mathcal{D} \begin{bmatrix} n+1, n+2 \\ 1, n+1 \end{bmatrix} \mathcal{D} \begin{bmatrix} 1, n+1, n+2 \\ 1, 2, n+2 \end{bmatrix} = 0.
\end{aligned} \tag{21}$$

Eq.(21) holds using a following Pfaffian identity. For any determinant A of an arbitrary square matrix

$$A \begin{bmatrix} a, b \\ d, e \end{bmatrix} A \begin{bmatrix} c \\ e \end{bmatrix} + A \begin{bmatrix} b, c \\ d, e \end{bmatrix} A \begin{bmatrix} a \\ e \end{bmatrix} + A \begin{bmatrix} c, a \\ d, e \end{bmatrix} A \begin{bmatrix} b \\ e \end{bmatrix} = 0, \tag{22}$$

where $a > b > c$. Leaving the proof of this identity later on we accept it.

Setting $A = \mathcal{D} \begin{bmatrix} 1 \\ n+2 \end{bmatrix}$, the first two terms of Eq.(21) become

$$\begin{aligned}
& A \begin{bmatrix} 2, n+1 \\ 1, n+1 \end{bmatrix} A \begin{bmatrix} n+2 \\ n+1 \end{bmatrix} - A \begin{bmatrix} 2, n+2 \\ 1, n+1 \end{bmatrix} A \begin{bmatrix} n+1 \\ n+1 \end{bmatrix} = -A \begin{bmatrix} n+2, n+2 \\ 1, n+1 \end{bmatrix} A \begin{bmatrix} 2 \\ n+1 \end{bmatrix} \\
& = -\mathcal{D} \begin{bmatrix} 1, 2 \\ n+1, n+2 \end{bmatrix} \mathcal{D} \begin{bmatrix} 1, n+1, n+2 \\ 1, n+1, n+2 \end{bmatrix}.
\end{aligned} \tag{23}$$

Similarly the last two terms of Eq.(21) are, by setting $A = \mathcal{D} \begin{bmatrix} n+2 \\ 1 \end{bmatrix}$, as

$$\mathcal{D} \begin{bmatrix} n+1, n+2 \\ 1, 2 \end{bmatrix} \mathcal{D} \begin{bmatrix} 1, n+1, n+2 \\ 1, n+1, n+2 \end{bmatrix}. \tag{24}$$

The sum of above two equations vanishes since

$$\mathcal{D} \begin{bmatrix} 1, 2 \\ n+1, n+2 \end{bmatrix} = \mathcal{D} \begin{bmatrix} n+1, n+2 \\ 1, 2 \end{bmatrix}, \tag{25}$$

which is one of the properties of $\mathcal{D}$ in Eq.(14). Eq.(20) is also proved analogously to the above.

The crucial identity Eq.(22) we have used is proved as follows. Let us consider a determinant of $n \times n$ anti-symmetric matrix A whose (i, j) element is $a_{ij} (= -a_{ji})$. Pfaffian $(1, 2, \dots, n)$ is defined by

$$\det A \equiv \begin{cases} (1, 2, \dots, n)^2 & \text{for even } n \\ 0 & \text{for odd } n \end{cases} \tag{26}$$

with $a_{ij} = (i, j)$.

By definition Pfaffian is expanded as

$$(1, 2, \dots, n) = \sum_{j=1}^n (-)^j (1, j)(2, 3, \dots, \hat{j}, \dots, n) \quad (27)$$

where hatted component is removed one.

We can also express a determinant of $(n+2)$ -th order matrix as Pfaffian of $2(n+2)$ -th order by

$$\det a_{ij} = (1, 2, \dots, n+1, n+2, n+2^*, n+1^*, \dots, 2^*, 1^*) \quad (i, j = 1, 2, \dots, n+2) \quad (28)$$

supplemented by the condition, $(i, j) = (i^*, j^*) = 0$, $(i, j^*) \equiv a_{ij}$.

From the antisymmetry property of Pfaffians it follows

$$\det a_{ij} = (-)^P (a, b, c, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, \hat{c}, \dots, n+2, n+2^*, \dots, \hat{d}^*, \dots, \hat{e}^*, \dots, 1^*, d^*, e^*), \quad (29)$$

where $(-)^P$ is ± 1 depending on the signature of permutation P of reordering. In terms of Pfaffians each terms of Eq.(22) are expressed as follows.

$$\begin{aligned} A \begin{bmatrix} a, b \\ d, e \end{bmatrix} &= (c, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, \hat{c}, \dots, \hat{d}^*, \dots, \hat{e}^*, \dots, 2^*, 1^*) \\ A \begin{bmatrix} c \\ e \end{bmatrix} &= (a, b, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, \hat{c}, \dots, \hat{d}^*, \dots, \hat{e}^*, \dots, 2^*, 1^*, d^*) \\ A \begin{bmatrix} b, c \\ d, e \end{bmatrix} &= (a, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, \hat{c}, \dots, \hat{d}^*, \dots, \hat{e}^*, \dots, 2^*, 1^*) \\ A \begin{bmatrix} a \\ e \end{bmatrix} &= (b, c, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, \hat{c}, \dots, \hat{d}^*, \dots, \hat{e}^*, \dots, 2^*, 1^*, d^*) \\ A \begin{bmatrix} a, c \\ d, e \end{bmatrix} &= (b, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, \hat{c}, \dots, \hat{d}^*, \dots, \hat{e}^*, \dots, 2^*, 1^*) \\ A \begin{bmatrix} b \\ e \end{bmatrix} &= (a, c, 1, \dots, \hat{a}, \dots, \hat{b}, \dots, \hat{c}, \dots, \hat{d}^*, \dots, \hat{e}^*, \dots, 2^*, 1^*, d^*) \end{aligned} \quad (30)$$

up to common signature factor $(-)^P$ provided that the order of a, b and c is kept in mind as in Eq.(22). Eq.(22) is expressed symbolically in terms of Maya diagram [4] as

$$\begin{aligned} & A \begin{bmatrix} a, b \\ d, e \end{bmatrix} A \begin{bmatrix} c \\ e \end{bmatrix} + A \begin{bmatrix} b, c \\ d, e \end{bmatrix} A \begin{bmatrix} a \\ e \end{bmatrix} - A \begin{bmatrix} a, c \\ d, e \end{bmatrix} A \begin{bmatrix} b \\ e \end{bmatrix} \\ &= \begin{array}{|c|c|c|c|} \hline a & b & c & d^* \\ \hline X & X & O & X \\ \hline \end{array} \times \begin{array}{|c|c|c|c|} \hline a & b & c & d^* \\ \hline O & O & X & O \\ \hline \end{array} \\ &+ \begin{array}{|c|c|c|c|} \hline O & X & X & X \\ \hline \end{array} \times \begin{array}{|c|c|c|c|} \hline X & O & O & O \\ \hline \end{array} \\ &- \begin{array}{|c|c|c|c|} \hline X & O & X & X \\ \hline \end{array} \times \begin{array}{|c|c|c|c|} \hline O & X & O & O \\ \hline \end{array} \\ &= \begin{array}{|c|c|c|c|} \hline O & O & O & X \\ \hline \end{array} \times \begin{array}{|c|c|c|c|} \hline X & X & X & O \\ \hline \end{array}. \end{aligned} \quad (31)$$

The last equality holds by virtue of Pfaffian identity. Both $\begin{array}{|c|c|c|c|} \hline O & O & O & X \\ \hline \end{array}$ and $\begin{array}{|c|c|c|c|} \hline X & X & X & O \\ \hline \end{array}$ consist of different numbers with and without asterisks and vanish by their definition. It completes the proof of the identity Eq.(22).

Next we discuss the second set of decomposition equations Eqs.(8) and (9). In contrast to the previous case the explicit form of ψ in Eq.(12) is required. $g_n(f_n)$ is the determinant of matrix whose $(i, j) \binom{i-1, j-1}{}$ element is

$$\begin{aligned} L_+^{i-1} L_-^{j-1} \psi &= L_+^{i-1} L_-^{j-1} (px - iqy) \\ &= pW_{i+j-1}(x) + (-1)^j iqW_{i+j-1}(y), \end{aligned} \quad (32)$$

where

$$W_{n+1}(z) = (z^2 - 1) \frac{d}{dz} W_n(z) \quad \text{with} \quad W_1(z) = z. \quad (33)$$

In this paper we restrict ourselves to prove for a case of $q = 0$ and the general case is left for future. In case of $p = 1$ and $q = 0$, $\psi = x$ and g_n and f_n are real functions depending only on x . Explicit forms of g_n and f_n are

$$g_n = \det \begin{pmatrix} W_1 & W_2 & \dots & W_n \\ W_2 & W_3 & \dots & W_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ W_n & W_{n+1} & \dots & W_{2n-1} \end{pmatrix}, \quad f_n = \det \begin{pmatrix} W_3 & \dots & W_{n+1} \\ \vdots & \ddots & \vdots \\ W_{n+1} & \dots & W_{2n-1} \end{pmatrix}. \quad (34)$$

Using Eq.(33) we can evaluate the determinant,

$$\begin{aligned} g_n &= \det \begin{pmatrix} x & 1 & 0 & 0 & \dots & 0 \\ x^2 - 1 & 2x & 2 & 0 & \dots & 0 \\ 2x(x^2 - 1) & 6x^2 - 1 & 12x & 12 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots \end{pmatrix} \\ &= \frac{A_n}{2} (x^2 - 1)^{\frac{n(n-1)}{2}} \left((x+1)^n + (x-1)^n \right), \end{aligned} \quad (35)$$

where the coefficient A_n is

$$A_n = (n-1)^2 (n-2)^4 \dots 2^{2(n-1)}. \quad (36)$$

The equation (35) is proved by induction. For $n = 1$ and $n = 2$ cases Eq.(35) holds. Assuming it for $n = l-1$ and $n = l$ cases we prove it for the case of $n = l+1$. Applying Jacobi's formula (4) to g_n we obtain, corresponding to the Toda molecule equation in Eq.(2),

$$g_{l-1} g_{l+1} = (L_X^2 g_l) g_l - (L_X g_l)^2, \quad (37)$$

where $L_X \equiv (x^2 - 1)\partial_x$. Using the assumed forms for g_l and g_{l-1} we find the expected form for g_{l+1} ;

$$g_{l+1} = \frac{A_{l+1}}{2} (x^2 - 1)^{\frac{(l+1)l}{2}} \left((x+1)^{l+1} + (x-1)^{l+1} \right). \quad (38)$$

Here we have used an equality obtained from the definition of A_n in Eq.(36);

$$A_{n-1} A_{n+1} = n^2 A_n. \quad (39)$$

Thus Eq.(35) is proved for $n = l + 1$. Quite analogously f_n is shown to be

$$f_n = \frac{A_n}{2}(x^2 - 1)^{\frac{n(n-1)}{2}} \left((x+1)^n - (x-1)^n \right). \quad (40)$$

The second set of the decoupled equations Eqs.(8) and (9) are

$$\begin{aligned} F(g_n \cdot f_n) &= 0, \\ F(g_n \cdot g_n + f_n \cdot f_n) &= 0, \end{aligned} \quad (41)$$

and F becomes in $q = 0$ and $c_n = -2n^2$ case

$$Fa \cdot b = \frac{1}{x^2 - 1} \left((L_X^2 a)b + a(L_X^2 b) - 2(L_X a)(L_X b) \right) - 2n^2 ab.$$

It is straight forward to show g_n and f_n in Eqs.(35) and (40) satisfy Eq.(41).

In this paper we have discussed a proof of Nakamura's conjecture on Tomimatsu-Sato solutions. In the proof a set of Pfaffian identities played crucial roles. The first set of decoupled equations for Ernst equation Eqs.(6) and (7) has been shown analytically. Here the explicit form of ψ in Eq.(12) is not required. The second set of decoupled equations Eqs.(8) and (9) has also proved for the restricted case of $q = 0$. They are corresponding to the (extended) Weyl solutions. There remains to show the most general case, $q \neq 0$. We expect that Pfaffian identities will play important roles in the analytic proof as in the case shown here. It is left as future problem.

References

- [1] R. Ward, Phil. Trans. R. Soc. Lond. **A315** (1985) 451
- [2] C. J. Papachristou and B. K. Harrison, Phys. Lett. **A190** (1994) 407,
V. A. Belinsky and V. E. Zakharov, Sov. Phys. JETP. **48** (1978) 985, **50** (1979) 1,
D. Mason, Phys. Rev. Lett. **41** (1978) 521.
- [3] R. Hirota, Phys. Rev. Lett. **27** (1971) 1192, R. Hirota, "Mathematical Theory of Solitons by Direct Method (Iwanami Pub. 1992; in Japanese).
- [4] M. Sato, Publ. RIMS Kokyuroku Kyoto Univ. **439** (1981) 30.
- [5] A. Nakamura, J. Phys. Soc. Japan. **62** (1993) 368.
- [6] R. Hirota, Y.Ohta and J. Satsuma, Prog. Theor. Phys. Supp. **94** (1988) 59.
- [7] A. Nakamura and Y. Ohta, J.Phys. Soc. Japan. **60** (1991) 1835